

## What is a Catchment Area?

- "The catchment area must be defined and justified by the [center] institution, based on the geographic area it serves."
- "It must be population based, e.g. using census tracts, zip codes, county or state lines, or other geographically defined boundaries."
- "It must include the local area surrounding the cancer center."
- "Cancer Centers occupy a unique role in their communities. They are expected to perform research relevant to their catchment area and engage the populations within their catchment area in the research they conduct and other Center activities."

NCI, CCSG FOA

The James

l

Geographically defined boundaries
 Zip code, county, state, etc.

How to Select a Catchment Area?

Primary service area – where clinics are located Market share – where patients come from

Institutional region - historic or alignment

Clinical and research geographic areas – where you have a footprint

Zip code, county, state, etc.

You decide

Options:

Rationale for decision needed

The James


THE ORDO STATE UNIVERSITY


## How to Characterize Your Catchment Area


- Who is in your catchment area?
  - Demographics
  - Special populations
- What is the cancer burden like?
- Risk factors
- Incidence
- Mortality
- Stage of diagnosis
- Survival


The James


## Information to Characterize the Catchment Area Information th

























| Risk Factors and Screening in Ohio  | WHITES          | BLACKS | APPALACHIANS |
|---|-----------------|--------|--------------|
| Current tobacco use, adults   | 22.8%           | 25.6%  | 30.8%        |
| Current tobacco use, teens  | 23.5%           | 16.0%  | 21.7%        |
| Obese adults age 18+  | 30.0%           | 34.5%  | 30.2%        |
| Ever had flex sig/colonoscopy, age 50+  | 65.2%           | 64.8%  | 62.9%        |
| Mammogram last 2 years, age 50-74   | 76.1%           | 86.4%  | 81.5%        |
| Pap smear, last 3 years, age 19-<65   | 77.2%           | 85.4%  | 81.1%        |
| To eliminate health disparities in Ohi<br>•High prevalence of obesity, especiall<br>•High prevalence of current tobacco ( | ly among blacks |        |              |

















## Thank You The James The James The James The James